

Evaluating the effect of dietary soybean oil on growth performances of juvenile severum (*Heros severus*)

Vesal S.E.^{1*}; Vosooghi A.R.²

Received: April 2016

Accepted: July 2016

Abstract

Evaluating the effect of dietary soybean oil (S.O) on growth performances of juvenile ornamental fish Severum (*Heros severus*) was the main objective of this study. To achieve this objective, 462 fish were considered which were divided into 7 test groups including 6 groups of fishes (with an initial average weight of 0.6 ± 0.05 g and length of 2.3 ± 0.35 cm) fed diets containing 6 levels of S.O (D₁(3), D₂(5), D₃(8), D₄(10), D₅(12), D₆(14)%) and the control group (CG) (receiving an oil-free diet) for 12 weeks with biometry intervals of 15 days. At the end of the test period, growth performance and nutritional parameters were measured. The study showed that the maximum weight was in fish fed diet containing 8% of S.O and also the maximum specific growth rate (SGR) (D₃, $p < 0.05$) was obtained for this group. The highest rate of daily weight gain (DWG) was obtained for D₃ which was significantly ($p < 0.05$) much more than that for D₄, D₆ and D₅ (the lowest increase). The highest length gain (LG) was observed in D₃ which was significantly more ($p < 0.05$) in comparison with the other treatments tested. The survival rate was over 80% in all test groups and showed no significant difference ($p > 0.05$). At the end of the period, fish were evaluated for estimated analysis of the tissue. The results of this study which was focused on the growth and development of ornamental fish *H. severus* fed with different levels of soybean oil suggest that the optimal level of this type of oil in their diet is 8% based on the growth parameters.

Keyword: Soybean oil, Lipid, Growth performances, Diet formulation, Severum (*Heros Severus*)

1- School of Marine Science and Technology, North Tehran Branch, Islamic Azad University (IAU), Tehran, Iran, Shahid Fallahi (Zaferaniyeh) Avenue, Postal code: 1987973133

2-Department of Aquaculture, Faculty of Fisheries, North Tehran Branch, Islamic Azad University (IAU), Tehran, Iran, Shahid Fallahi (Zaferaniyeh) Avenue, Postal code: 1987973133

*Corresponding author's Email: Ehsan.Vesal.64@gmail.com

Introduction

Nowadays, in addition to edible fish culture, breeding of different types of ornamental fish is taken into consideration and research projects are conducted enthusiastically in this field. The economic importance of ornamental fish is not lower than that of edible fish. In this regard, an important aspect of aquaculture is nutrition that fish farmers should pay special attention to, since a large part of the fish production costs is allocated to this part (Firouzbaksh *et al.*, 2011). However, little information is available on the nutritional requirements of ornamental fish (Kruger *et al.*, 2001; Miller and Mitchell., 2008; Güroy *et al.*, 2012).

The reproduction and breeding industry of ornamental fish is developing in line with the growth of aquaculture industry in the world. More than 1000 types of freshwater fish, which consist of 100 families, are commercial ornamental fish (FAO, 2003). In this regard, this field of fisheries science has become a large industry and a profitable business. On the other hand, health and nutrition of ornamental fish are two decisive factors in the ornamental fish industry (Ghosh *et al.*, 2008). Thus, food has a key role in health, appropriate natural behavior, improving the phenotype, the color of ornamental fish and also their growth, development and reproduction (Erdogan *et al.*, 2012). Also, the apparent growth is affected by the amount of food intake and food

absorption into the body (Buurm and Diana, 1994).

In this regard, formulated commercial diets for salmon often consist of 40-50% crude protein and 12-17% crude lipid that are used in ornamental fish farms regardless of nutritional behavior and ecology of the natural nutrition of this fish. As a result, it leads to decreasing growth performance, and economically, it is not affordable (Royes *et al.*, 2006).

Trading volume index of ornamental fish is about one billion dollars in the world (Erdogan *et al.*, 2012). The global supply of vegetable oils clearly represents the largest source of lipid for animal diets (Soyatech, 2009). Soybean oil (S.O), as a source of fat in the diet, is used in published studies on fish such as sturgeon (*Acipenser schrencki*) (Xiao and Cheng, 2001), common carp (*Cyprinus carpio*) (Fontagne *et al.*, 2000), Japanese flounders (Lee *et al.*, 2000) (*Paralichthys olivaceus*). The use of S.O in the above studies cannot indicate practical and reliable application of this fat source, and S.O should be evaluated based upon target species (Turchini *et al.*, 2011).

S.O has about 60 g of EFA /100 g of fatty acids and EFA in S.O is dominated by 18:2n-6 group (Tacon, 1990). S.O has been considered among available vegetable oils because of its availability, low cost and high amounts of essential fatty acids (Bell *et al.*, 2001; Caballero *et al.*, 2003). Grisdale-Helland *et al.* (2001) examined the soybean oil diet on growth performance

and feed in Atlantic trout (*Salmo salar*). They found that fish fed the diet containing 27.8% of S.O is in the second rank of growth among the diets tested, and they did not show any significant differences in different test groups (containing 27.8% fish oil and in a 1:1 ratio for fish oil and soybean oil each one 13.9%). Ng *et al.* (2012) did not observe any significant differences in their study on growth parameters in red hybrid tilapia fish (*Oreochromis* sp.) fed diets containing vegetable oils (soybean, palm and flax) and fish oil. Jiang *et al.* (2012) examined the effects of diets containing soybean oil and fish oil on the growth of catfish (*Pelteobagrus vachelli*). In their experiment, they found that diets containing 8% S.O did not show any significant differences in weight gain (WG) during the 80 days of feeding. Lochman and Brown (1997) examined the average weight gain in goldfish (*Carassius auratus*) fed with a diet containing 4% fish oil (cod liver oil) and S.O (4%) for a period of 42 days. They found that there was no significant difference on weight gain in gold fish fed with a diet containing 4% S.O throughout this period. Regost *et al.* (2003) studied the growth performance of Turbot (*Psetta maxima*) fed with diets containing 9% S.O and 9% fish oil, and observed a significant difference in final weight and growth rate of the fish. Deng *et al.* (2013) examined the growth factors of Pacific threadfin (*Polydactylus sexfilis*) fed a

diet containing fish oil, S.O and in a 1:1 ratio for fish oil and S.O.

Their results suggested that diets containing 8% S.O should provide adequate levels of essential fatty acids to support normal growth of juvenile Pacific threadfin under the current feeding conditions.

The purpose of this research is to produce proper diets with emphasis on optimum levels of fat using S.O for ornamental species of *severus* (*Heros severus*) according to the growth parameters in the aquarium environment.

Materials and methods

Experimental diets

In this study, 6 experimental diets of formulated isonitrogenous with different levels of S.O (3, 5, 8, 10, 12, 14%) and a control group (experimental diet without oil) were prepared. Diets with raw material (Table 1) were prepared in accordance with the formula written by software 2.8 Win Feed (Cambridge, UK). The raw materials were ground (diameter 1 mm) using an electric mill. According to the provided formula the level of each one was calculated and weighed. Then, these materials were mixed to form a paste by adding water. The amount of oil needed was weighed according to the formula and then added to the paste mixture.

Table 1: Ingredient and proximate composition of experimental diets.

Ingredient (g/10g)	Experimental diets						
	D ₁ (3)	D ₂ (5)	D ₃ (8)	D ₄ (10)	D ₅ (12)	D ₆ (14)	CG
Fish meal	1.89	1.9	1.93	1.94	1.82	1.68	1.91
Wheat meal	1.43	1.35	1.22	1.14	0.75	0.27	1.52
Wheat gluten	2.02	2.05	2.09	2.12	2.12	2.12	1.99
Soybean meal	1.75	1.71	1.65	1.61	1.9	2.28	1.75
Oil	0.15	0.36	0.67	0.88	1.11	1.34	-
Vitamin premix	0.2	0.2	0.2	0.2	0.2	0.2	0.2
Mineral premix	0.15	0.15	0.15	0.15	0.15	0.15	0.15
*Additives	1.05	1.05	1.05	1.05	1.05	1.05	1.05
Dicalcium phosphate	1.36	1.23	1.04	0.91	0.9	0.91	1.42
	Proximate analyses (% DM*)						
Dry matter	90.88	91.11	91.51	91.85	92.04	92.38	89.62
Crude protein	44.42	44.80	45.30	45.80	46.20	46.83	43
Crude fat	6.31	7.70	10.22	12.80	15.01	16.55	4.59
Ash	14.70	14.85	13.62	12.50	12.75	13.10	17

*Additives : Astaxanthin 3%, antioxidants 0.1%, binder3%, moderate inhibitor 0.4%, D L methionine 1%, lysine 1%, Garlic Powder 2%; *DM, dry matter.

The paste was passed through a meat grinder with a pore plate with 2 mm pores (in diameter) and the resulting paste strings were placed in aluminum trays. After that, trays were kept in an oven at a temperature of 65°C for 8 hours, and then dried under the hood in sterile conditions. Finally, these threads were cut by the cutter according to the size of the fish's mouth, sealed in plastic bags and kept at 4°C until usage.

Design of experiment

In this study, 462 numbers of Severum with an average weight of 0.6±0.05 g and an average length of 2.3±0.35 cm were obtained (all from the same parent). After preparation of 21 aquaria with dimensions of 30×50×70 cm and volume of 100 liters, the aquaria were equipped with an aeration system and heater to maintain the water temperature. Fish were put into 7

experimental groups; 6 groups with diets containing S.O at levels of 3 (D₁), 5 (D₂), 8 (D₃), 10 (D₄), 12 (D₅), 14 (D₆) and 1 control group (CG) (oil-free diet) for 12 weeks; each one was replicated 3 times. Fish were stocked randomly inside the aquaria (22 per aquarium) and treatments and group numbers were assigned to aquaria according to the type of food. For adaptation, fish were fed a basic diet and were kept in the conditions of new aquaria for 8 days. To protect water quality during the period, bio-filters were used and 50% of water was replaced every three days. The dissolved oxygen and pH of the water were measured every two days using specialized test kits via the colorimetric method. Fish were fed 3 times a day (08:00, 11:00 and 16:00 hours) on the basis of biomass obtained in each biometric assessment, and the amount of food was calculated

according to appetite of fish and based upon 4% of the fish weight. The main stage of the test started and continued for 84 days after completion of the adaptation period, and biometric assessments were conducted once every 15 days. To measure weight and length, a scale with a precision of 0.01 g and a precision ruler were used, respectively. Feedings were stopped for 15 hours before and after biometric assessment to reduce fish stress in the course of the procedure.

Calculation of growth indices, feed indices and Survival rate

At this stage, according to data obtained from biometric assessments, growth parameters which include specific growth rate, the percentage of body weight gain, the percentage of body length gain, survival rate of average daily gain, condition factors and nutritional indices including optional food intake and feed conversion ratio were measured as follows (Austreng, 1978; De Silva and Anderson, 1995; Yanbo and Zirong, 2006; Gümüş, 2011).

Specific growth rate (%); $SGR_w = [\ln \text{ final body weight} - \ln \text{ initial body weight} / \text{days of experiment}] \times 100$.

Feed conversion ratio; $FCR = \text{Feed intake (g)} / \text{weight gain (g)}$.

Weight gain; $WG (g) = (\text{Final weight} - \text{initial weight}) / (\text{initial weight}) \times 100$.

Length gain; $LG (mm) = (\text{Final length} - \text{initial length}) / (\text{initial length}) \times 100$.

Condition factor; $CF = (\text{Final weight (g)} / (\text{fork length (cm)})^3) \times 100$.

Daily growth rate; $DGR (g) = ((\text{Final body weight}) - (\text{initial body weight}) / \text{initial body weight} \times \text{days})$.

Voluntary feed intake; $VFI = \text{Dry feed intake} / [(\text{initial} + \text{final fish biomass})/2]$

Specific growth rate (%); $SGR_L = [\ln \text{ final body length} - \ln \text{ initial body length} / \text{days of experiment}] \times 100$

Survival rate (%); $SR = (\text{Final fish number} / \text{Initial fish number}) \times 100$.

Carcass analysis

At the end of the feeding trial, 8 fish were randomly selected from treatments fed with diets consisting the levels of 3(D₁), 5(D₂), 8(D₃), 10(D₄), 12(D₅), 14(D₆) oil content and the control group (without oil) (CG), and transferred to the laboratory. The total carcass protein content of body and crude oil were measured using the Kjeldahl method and Soxhlet method according to the instructions of AOAC (1990), respectively. Moisture was also measured by the method described in AOAC (1990) (Table 3).

Statistical analysis

The mean values related to different groups of data were calculated in Excel and then imported into SPSS version 20 to be checked for the significant differences between results related to different treatments. Normal scattering of data was determined using the Kolmogorov-Smirnov test, and then the significant differences between treatments were investigated using One-Way ANOVA. After observation of significant difference, post hoc LSD in

the ($p < 0.05$) level was used for investigation of significant differences between the groups.

Results

The parameters of water quality were maintained as follows: water temperature = 28.2°C, pH = 6.8 to 7.8, dissolved oxygen: 6.8 to 8.3 ppm, nitrite = 0.02 mgL⁻¹, and total hardness: 215 mgL⁻¹. Growth indices, feed conversion ratio and the percentage of survival in severum (*H. severus*) fed with diets containing different levels of S.O are shown in Table 2.

The percentage of body weight gain (WG), which is a function of final weight, was significantly different ($p < 0.05$) among the studied groups at the end of the period. In fact, the highest weight gain was recorded in fish fed diets containing 8% oil (D₃) in that the specific growth rate of weight (SGR_W) of this group was considerably higher than the studied treatments at the end of the period ($p < 0.05$). However, the lowest weight gain was in fish fed diets containing 12% oil (D₅). The daily weight gain rate (DGR) was calculated in fish of different treatments and found that the maximum value was in fish fed a diet containing 8% oil, which showed significant differences ($p < 0.05$) with treatments D₄, D₅ and D₆, and the lowest value was in fish fed a diet containing 12% oil (D₅).

Feed conversion ratio (FCR) of fish fed diets containing 8 and 10% oil was the lowest at the end of the period. Also, highest FCR was related to fish

fed the diet containing 12% oil (D₅). At the end of the experiment, survival rate (SR) in all treatments had no significant differences, and was more than 80%. Also, no sign of disease was observed in all fed fish ($p > 0.05$). With statistical analysis and study of Table 2 in terms of specific growth rate of length (SGR_L), fish fed a diet containing 8% oil (D₃) had higher SGR_L than the rest of treatments at the end of the period ($p < 0.05$). The lowest SGR_L was in fish fed a diet containing 12% oil (D₅). Condition factor (CF) of fish of different treatments was calculated and it was found that the maximum amount was for fish fed a diet containing 10% oil (D₄), which was significantly different ($p < 0.05$) with that in other treatments and the lowest was for fish fed the diet of 12% oil (D₅). The highest and lowest amounts of voluntary food intake (VFI) were in fish fed diets of 12% oil (D₅) and 8% oil (D₃), respectively. Fish fed diets of 8% oil (D₃) showed the highest growth that had significant differences with that in the other treatments ($p < 0.05$), while fish fed a diet containing 12% oil showed the lowest growth rates (Table 2).

Table 2: Growth parameters of *Heros severus* fed the experimental diets containing soybean oil for 84 days.

Growth parameters	3 (%)	5(%)	8(%)	10(%)	12(%)	14(%)	CG (%)
FCR (%)	1.69±0.1 ^b	1.55±0.05 ^d	1.53±0.11 ^c	1.53±0.11 ^c	2.5±0.41 ^a	1.69±0.1 ^b	1.64±0.06 ^e
SGR _w (%)	17.48±1.97 ^b	18.7±0.64 ^a	19.39±2.42 ^a	16.5±2.18 ^b	9.51±3.17 ^d	14.59±0.96 ^c	15.53±1.73 ^c
SGR _L (%)	54.88±12.62 ^b	56.26±4.68 ^a	58.54±21.38 ^c	53.41±12.27 ^c	49.62±29.4 ^e	54.65±13.19 ^d	42.19±24.02 ^f
WG (g)	14.33±1.61 ^b	15.3±0.52 ^a	15.9±1.98 ^a	13.53±1.79 ^c	7.80±2.6 ^e	11.96±0.38 ^d	13.18±0.84 ^c
LG (mm)	45±10.34 ^a	46.13±3.84 ^a	48±17.53 ^b	43.8±10.06 ^b	40.68±24.11 ^d	44.81±10.81 ^c	34.64±11.54 ^f
SR (%)	100±0.00 ^a	100±0.00 ^a	93.33±5.43 ^a	100±0.00 ^a	80±9.42 ^b	93.33±5.43 ^a	100±0.00 ^a
CF (%)	15.61±0.45 ^b	15.62±0.88 ^b	14.49±3.62 ^c	15.94±0.71 ^a	11.22±3.55 ^e	13.47±2.18 ^d	13.21±3.24 ^d
VFI	69.25±4.31 ^b	63.59±2.09 ^c	62.88±4.9 ^c	63.12±4.84 ^c	102.55±16.98 ^a	69.33±4.11 ^b	62.62±2.81 ^c
DGR (gr)	69.56±1.32 ^a	70.48±0.33 ^a	70.75±1.31 ^a	68.9±1.37 ^b	59.83±4.48 ^e	67.8±0.38 ^b	68.27±0.81 ^b

* Different superscript in each row indicate significant differences among treatments ($p < 0.05$).

Table 3: Proximate composition (%) of whole body of *Heros severus* fed diets containing soybean oil.

Diet	Experimental diets (%)						
	D1(3)	D2 (5)	D3 (8)	D4 (10)	D5 (12)	D6 (14)	CG
Crude protein	68.1±0.18	67.30±0.17	65.52±0.32	62.70±0.34	62.80±0.33	63.48±0.27	65.1±0.38
Crude lipid	9.7±0.31	13.95±0.4	20.50±0.36	34.69±0.29	17.50±0.31	23.70±0.43	8.9±0.25
Moisture	24.3±0.14	25.60±0.33	25.50±0.36	25.70±0.32	25.50±0.32	27.65±0.8	25.1±0.52
Ash	6.45±0.15	6.70±0.09	6.90±0.14	7.20±0.13	7.40±0.34	7.50±0.35	7.70±0.33

Discussion

This research was conducted to determine the optimum level of lipid using S.O for growth and maintenance of ornamental severum (*H. severus*). The results show that infants of severum fed the 8% oil diet had higher growth rates than the other levels. The use of oil in the diet of fish such as energy sources in diets of fish had effects on growth, survival and body composition (Sener and Yildiz, 2003; Almaida-Pagan *et al.*, 2007).

This is the first study on the impact of diets containing different lipid levels based upon S.O on growth and food productivity and body composition of *H. severus* ornamental species. The results of this study suggest that fish fed diets containing 8% oil (D₃) with a higher weight gain showed better specific growth rates than other groups.

It indicates that nutrients available in the diet D₃ provide the nutritional requirements needed for faster growth of severum, compared to other groups. It means that increase in the amount of oil and energy in the diet of treatment (D₃) decreases protein of tissue and raises fat. In other words, it increases the fat of tissue. Piedecausa *et al.* (2007) reported that the highest level of fatty acids, 18: 2n-6 in the S.O can increase fat storage in the fish body, and also Ng *et al.* (2003) claimed that a high content of fatty acids in the diet leads to increasing the storage of these acids in muscle tissue. Hence, we can say that there is a positive association between dietary fat and fat content of total body for many fish species (Chou and Shiau, 1996; Du *et al.*, 2005; El-Marakby, 2006; Martins *et al.*, 2007).

Fish fed with diets of 12% (D₅) showed the lowest SGR which indicates that the diet did not have the adequate fat for growth and also there is an imbalance between fat and protein content. In the present study, the maximum value of CGR was in fish fed diets of 8% oil (D₃) that is similar to the study which was carried out by Deng *et al.* (2013) on Pacific threadfin (*P. sexfilis*). The highest SGR was obtained in S.O 8%. Since SGR decreases with increasing fish size, lower values observed for specific growth rate (SGR) in some studies can be explained by the evolution of fish (Sunde *et al.*, 1998).

Peng *et al.* (2008) reported the amount of weight gain for young fish of black marine wire (*Acanthopagrus schlegeli*) fed diets of 9% S.O was 2.20 g. In this study, the best FCR for Severum was achieved using diets of 8 and 10% oil. The potential reason which can be considered for better growth of D₃ and D₄ is that in these are the best diet sources of nutrients to be consumed by fish. In this work, the highest FCR was for fish fed with the diet of 12% S.O which was similar to the results obtained in the investigation of Ricardo *et al.* (2001) on the performance and composition of fatty acids of fish Surubim (*Pseudoplatystoma coruscans*) fed with vegetable oils and animal.

The highest FCR was observed in treatments fed a diet containing 12% S.O and other vegetable oils tested (Corn oil and Linseed oil). In this research, the percentage of weight gain

rises along with the increase of fat to the level of 8%. Subhadra *et al.* (2006) suggested that changes in fish growth can be the result of a combination of diets and the length of feeding period. In this study, by increasing oil level from 8 to 14%, weight loss was observed and growth factors had no significant differences. So, we can say that the use of this amount of S.O in the diet of *H. severus* is not desirable. Lachman and Brown (1997) in their study found that the average weight gain in goldfish (*Carasius auratus L.*) fed a diet containing 4% S.O for a period of six weeks cannot show a significant difference. In this study, the largest percentage of length gain (LG) was in fish fed diets of 8% oil. By contrast, Jiang *et al.* (2012) found that catfish (*P. vachelli*) fed diets containing different levels of S.O did not show any significant differences in growth parameters (weight and height) after 80 days of test period. In this work, the condition factor was higher in fish fed diets of 10% oil (D₄) in comparison with the other treatments. The cause of high CF in this treatment could be because of the lower amount of protein and higher amount of fat in the body of Severum fish which was involved. As a result, by increasing body weight, because of the lower effect of protein on bone growth, it did not have good growth (Vesal *et al.*, 2016). In this experiment, CF in fish fed the diet of 12% S.O (D₅) was lower compared to other treatments. Deng *et al.* (2013) found that the CF was the lowest for

pacific threadfin fish (*P. sexfilis*) fed diets containing 8% S.O which showed significant differences that were not in accordance with the findings of this research. Regost *et al.* (2003) in their study observed a significant difference between the final weight and growth rate of Turbot fish (*P. maxima*) fed diets of S.O 9%. They found that the treatments fed diets containing vegetable oils cannot only have a significant impact on the growth of the fish. In this study, the maximum amount of voluntary food intake (VFI) was in fish fed the diet containing 12% oil (D₅) which had the lowest growth rate and the highest FCR.

Peng *et al.* (2014) investigated the growth parameters in the Turbot fish (*Scophthalmus maximus* L.) by replacing the different levels of fish oil with S.O, and found that the use of 7.5% S.O in the diet had an intermediate effect on food efficiency and food consumption of this fish, compared to other diets tested (mixed 1:3 fish oil+2:3 S.O and vice versa). Also, fish fed the lowest level of S.O had the highest food intake (FI) in comparison with the other groups ($p < 0.05$).

Previous studies have shown that fish are able to adjust their food consumption based upon diet quality using a self-feeding device when the time of food availability is not long enough. Accordingly, feeding with a diet, which is faced with a weak adoption, has often been rejected and we can say that this nutritional behavior

of fish is responsible for the amount of uneaten food (Vesal *et al.*, 2016).

There were no signs of disease among fish. However, the lowest survival rate was obtained in the diet containing 12% oil (D₅). But, due to very little differences, it seems that the mortality rate is independent of the fat level in the diet. It has been reported that the use of S.O in the diet of *A. schlegeli* cannot have any effect on the survival of this species (Peng *et al.*, 2008).

In this study, the efficiency growth of *H. severus* fed different levels of S.O in the diet showed that the optimal level of fat for this species is using 8% of this oil. The present investigation offers a preliminary approximation of the lipid requirements of *H. severus* regarding the use of S.O. Nonetheless, according to the current results, the quantitative necessities of essential fatty acids have not been evaluated and require further investigations.

References

- Almaida-Pagan, P.F., Hernandez, M.D., Garcia, B., Madrid, J.A., De Costa, J. and Mendiola, P., 2007. Effects of total replacement of fish oil by vegetable oils on n-3 and n-6 polyunsaturated fatty acid desaturation and elongation in sharpsnout seabream (*Diplodus puntazzo*) hepatocytes and enterocytes. *Aquaculture*, 272, 589–598.

- AOAC, 1990.** Official methods of analysis, 14th edition. Association of Official Analytical Chemists, Arlington, VA., 1102P.
- Austreng, E., 1978.** Digestibility determination in fish using chromic oxide marking and analysis of contents from different segments of the gastrointestinal tract. *Aquaculture*, 13, 265-272.
- Bell, J.G., McEvoy, J., Tocher, D.R., McGhee, F., Campbell, P.J. and Sargent, J.R., 2001.** Replacement of fish oil with rape seed oil in diets of Atlantic salmon (*Salmo salar*) affects tissue lipid compositions and hepatocyte fatty acid metabolism. *Journal of Nutrition*, 131, 1535–1543.
- Buurm, B.J. and Diana, J.S., 1994.** Effects of feeding frequency and handling on growth and mortality of cultured walking cat fish *clarius fuscus*. *Journal of the World Aquaculture Society*, 25, 175-182.
- Caballero, M.J., Izquierdo, M.S., Kjørsvik, E., Montero, D., Socorro, J., Fernandez, A.J. and Rosenlund, G., 2003.** Morphological aspects of intestinal cells from gilthead sea bream (*Sparus aurata*) fed diets containing different lipid sources. *Aquaculture*, 225, 325–340.
- Chou, B.S. and Shiau, S.Y., 1996.** Optimal dietary lipid level for growth of juvenile hybrid tilapia, *Oreochromis niloticus* × *Oreochromis aureus*. *Aquaculture*, 143, 185–195.
- De Silva, S.S. and Anderson, T.A., 1995.** Fish nutrition in aquaculture. London, Chapman and Hall, 319P.
- Deng, D.F., Ju, Z.Y., Dominy, W.G., Conquest, L., Smiley, S. and Bechtel, P.J., 2013.** Effect of replacing and growth performance of juvenile Pacific threadfin (*Polydactylus sexfilis*). *Aquaculture*, 422–423, 91–97.
- Du, Z.Y., Liu, Y.J., Tian, L.X., Wang, J.T., Wang, Y. and Liang, G.Y., 2005.** Effect of dietary lipid level on growth, feed utilization and body composition by juvenile grass carp (*Ctenopharyngodon idella*). *Aquaculture Nutrition*, 11, 139–146.
- El-Marakby, H.I., 2006.** Effect of dietary sources and levels of lipids on growth performance and feed utilization of fry Nile tilapia, *Oreochromis niloticus* (L.) (*Teleostei:Perciformes*). *Journal of Fisheries and Aquatic Science*, 1, 117–125.
- Erdogan, F., Erdogan, M. and Gümüş, E., 2012.** Effects of dietary protein and lipid levels on growth performances of two African cichlids (*Pseudotropheus socolofi*) and (*Haplochromis ahli*), *Turkish Journal of Fisheries and Aquatic Sciences*, 12: 453-458.
- FAO, 2003.** Yearbook fishery statistic, *Aquaculture Production*, 96, 2.
- Firouzbakhsh, F., Noori, F., Khalesi, M.K. and Jani-Khalili, K., 2011.** Effects of a probiotic, protexin, on the growth performance and hematological parameters in the

- Oscar (*Astronotus ocellatus*) fingerlings. *Fish Physiology and Biochemistry*, Doi: 10.1007/s10695-011-9481-4.
- Fontagne, S., Burtaire, L., Corraze, G. and Bergot, P., 2000.** Effects of dietary medium-chain triacylglycerols (tricaprylin and tricaproin) in phospholipid supply on survival, growth and lipid metabolism in common carp (*Cyprinus carpio*) larvae. *Aquaculture*, 190, 289–303.
- Ghosh, S., Sinha, A. and Sahu, C., 2008.** Dietary probiotic supplementation on growth and health of live-bearing of ornamental fishes. *Aquacultura Nutrition*, 14, 289-299.
- Grisdale-Helland, B., Ruyter, B., Rosenlund, G., Obach, A., Helland, S.J., Sandberg, M.G., Standal, H. and Røsjø, C., 2001.** Influence of high contents of dietary soybean oil on growth, feed utilization, tissue fatty acid composition, heart histology and standard oxygen consumption of Atlantic salmon (*Salmo salar*) raised at two temperatures. *Aquaculture*, 207, 311 – 329.
- Gümüş, E., 2011.** Effect of replacement of fishmeal with sand smelt (*Atherina boyeri*) meal on growth, feed utilization and body composition of mirror carp fry (*Cyprinus carpio*). *Kafkas Üniversitesi Veteriner Fakültesi Dergisi*, 17(3), 363-369.
- Güroy, D., Şahin, I., Güroy, B., Altın, A. and Merrifield, D.L., 2012.** Effect of dietary protein level on growth performance and nitrogen excretion of the yellow tail cichlid, *Pseudotropheus acei*. *The Israeli Journal of Aquaculture*. Bamidgheh, IJA: 64.2012.684, 6P.
- Jiang, X., Chen, L., Qin, J., Qin, Ch., Jiang, H., Li. and Er., 2012.** Effects of dietary soybean oil inclusion to replace fish oil on growth, muscle fatty acid composition, and immune responses of juvenile darkbarbel catfish, *Pelteobagrus vachelli*. *African Journal of Agricultural Research*, 8(16), 1492-1499.
- Kruger, D.P., Britz, P.J. and Sales, J., 2001.** Influence of varying dietary protein content at three lipid concentrations on growth characteristics of juvenile word tails (*Xiphophorus helleri* Heckel, 1848). *Sciences and Conservation Aquarium*, 3(1), 275–280. Doi: 10.1023 / A : 1013150314719.
- Lee, S.M., Cho, S.H. and Kim, K.D., 2000.** Effects of dietary protein and energy levels on growth and body composition of juvenile flounder *Paralichthys olivaceus*. *Journal of the World Aquaculture Society*, 31, 306–315.
- Lochmann, R. and Brown, R., 1997.** Soybean-lecithin supplementation of practical diets for juvenile goldfish (*Carassius auratus*). *Journal of the American Oil Chemists' Society*, 74(2), 149-152.

- Martins, D.A., Valente, L.M.P. and Lall, S.P., 2007.** Effects of dietary lipid level on growth and lipid utilization by juvenile Atlantic halibut (*Hippoglossus hippoglossus*, L.). *Aquaculture*, 263, 150–158.
- Miller, S.M. and Mitchell, M.A., 2008.** Ornamental fish. In: M. Mitchell and T. Tully (Eds.). *Manual of Exotic Pet Practice*. Saunders Publ., St. Louis: 39-72.
- Ng, H.H., Ciccone, D.N., Morshead, K.B., Oettinger, M.A. and Struhl, K., 2003.** Lysine 79 of histone H3 is hypomethylated at silenced loci in yeast and mammalian cells: a potential mechanism for position-effect variegation. *Proceedings of the National Academy of Sciences, USA* 100, 1820–1825.
- Ng, W.-K., Chong, C-Y., Wang, Y. and Romano, N., 2012.** Effects of dietary fish and vegetable oils on the growth, tissue fatty acid composition, oxidative stability and vitamin E content of red hybrid tilapia and efficacy of using fish oil finishing diets. *Aquaculture*, 372–375(2013), 97–110.
- Peng, M., Xu, W., Mai, Zhou, K.H., Zhang, Y., Liufu, Z., Zhang, K. and Ai, Q., 2014.** Growth performance, lipid deposition and hepatic lipid metabolism related gene expression in juvenile turbot (*Scophthalmus maximus* L.) fed diets with various fish oil substitution levels by soybean oil. *Aquaculture*, 433, 442–449.
- Peng, S., Chen, L., Qin, J.G., Hou, J., Yu, N., Long, Z., Ye, J. and Sun, X., 2008.** Effects of replacement of dietary fish oil by soybean oil on growth performance and liver biochemical composition in juvenile black seabream, *Acanthopagrus schlegelii*. *Aquaculture*, 276, 154–161.
- Piedecausa, M.A., Mazon, M.J., Garcia Garcia, B. and Hernandez, M.D., 2007.** Effects of total replacement of fish oil by vegetable oils in the diets of sharpsnout seabream (*Diplodus puntazzo*). *Aquaculture*, 263, 211-219.
- Regost, C., Arzel, J., Robin, J., Rosenlund, G. and Kaushik, S.J., 2003.** Total replacement of fish oil by soybean or linseed oil with a return to fish oil in turbot (*Psetta maxima*): 1. Growth performance, flesh fatty acid profile, and lipid metabolism. *Aquaculture*, 217, 465–482.
- Ricardo, C. M., Eurico, P. C., Leandro, P., Luiz, C. T., 2002.** Performance and fatty acid composition of surubim (*Pseudoplatystom couscans*) fed diets with animal and plant lipids. *Aquaculture* 209, 233 – 246
- Royes, J.A., Murie, D.J. and Francis-Floyd, R., 2006.** Effects of varying dietary protein and lipid levels on growth performance and hepatocyte changes in juvenile African cichlids. *The Journal of the World Aquaculture Society*, 37(1), 48-59.

- Sener, E. and Yildiz, M., 2003.** Effect of different oil on growth and body composition of rainbow trout (*Oncorhynchus mykiss* W., 1792) juveniles. *Turkish Journal of Fisheries and Aquatic Sciences*, 3, 111–116.
- Soyatech, 2009.** Soya and oilseed bluebook. Southwest Harbor, ME: Soyatech, pp. 324–368.
- Subhadra, B., Lochmann, R., Rawles, S. and Chen, R., 2006.** Effect of dietary lipid source on the growth, tissue composition and hematological parameters of largemouth bass (*Micropterus salmonides*). *Aquaculture*, 255, 210–220.
- Sunde, L.M.; Imsland, A.K. and Folkvord, A., 1998.** Effects of size grading on growth and survival of juvenile turbot at two temperatures. *Aquaculture International*, 6, 19-32.
- Tacon, A.G.J., 1990.** Standard methods for the nutrition and feeding of farmed fish and shrimp. Argent Laboratories Press, Redmond, Washington, USA. 128P.
- Turchini, G.M., Torstensen, B.E. and Ng, W.K., 2011.** Fish oil replacement in finfish nutrition. *Reviews in Aquaculture*, 1, 10–57.
- Vesal, S.E., Mooraki, N. and Vosooghi, A.R., 2016.** Effect of dietary Fish oil on growth responses of severum (*Heros severus*). *AACL Bioflux*, 9(1), 81-90.
- Xiao, Y. and Cheng, Y., 2001.** Effects of dietary lipid content on growth rate and liver lipids composition in *Acipenser schrencki*. *Journal of Oceanography*, 20, 376–38.
- Yanbo, W. and Zirong, X., 2006.** Effect of probiotic for common carp (*Cyprinu carpio*) based on growth performance and digestive enzymes activities. *Animal Feed Science and Technology*, 127, 283-292.